Can a beam splitter prevent correlation



Overview

The second-order anticorrelation on a beam splitter represents these phenomena where it cannot be achieved classically.

Key Takeaways

A beam splitter can induce anticorrelation between photons, but it does not universally prevent correlation; the effect depends on the quantum state and coherence of the input light.

How Beam Splitters Affect Correlation

A beam splitter is an optical device that divides an incoming light beam into transmitted and reflected components, often in a 50: 50 ratio. In classical optics, the outputs of a beam splitter can remain correlated depending on the coherence of the input beams. However, in quantum optics, the situation is more nuanced.

Quantum Anticorrelation

When single photons are incident on a beam splitter, quantum mechanics predicts anticorrelation: a single photon can only be detected in one output port at a time, never both simultaneously. This phenomenon, observed in experiments like the Hong-Ou-Mandel effect, arises from destructive quantum interference between the two possible paths of the photon. The anticorrelation is a direct consequence of the wave nature and quantum superposition of the input fields, not merely the particle nature of photons.

© 2026 DOLPHIN FIBER SYSTEMS - All rights reserved

Dependence on Input StatesThe ability of a beam splitter to prevent correlation depends on the statistical properties of the input light: Single-photon or antibunched states: Strong anticorrelation occurs, reducing coincidences at the outputs. Coherent or thermal light: Partial correlations may remain, and classical intensity correlations can still be observed. Entangled photons: Correlations can be manipulated via interference, but the beam splitter alone does not eliminate all correlations; it redistributes them according to quantum superposition.

Summary

A beam splitter does not inherently "prevent" correlation in all cases. Instead, it can create anticorrelation for certain quantum states, particularly single-photon or nonclassical states, due to interference effects. For classical or coherent light, correlations may persist, and the output statistics depend on the input coherence and photon statistics.

The phenomenon highlights the interplay between quantum superposition, coherence, and measurement in determining correlation outcomes at a beam splitter.

Article Content

The origin of anticorrelation for photon bunching on a beam splitter

The second-order anticorrelation on a beam splitter represents these phenomena where it cannot be achieved classically. Here, the anticorrelation of nonclassicality on a beam splitter is interpreted in a

The origin of anti-correlation for photon bunching on a beam splitter

The second-order anti-correlation on a beam splitter (BS) has been intensively studied over the last several decades due to its weird phenomenon of sub-Poisson statistics resulting in photon bunching.

Beam splitter

To reduce loss of light due to absorption by the reflective coating, so-called "Swiss-cheese" beam-splitter mirrors have been used. Originally, these were sheets of

Fundamental properties of beamsplitters in classical and quantum optics

A lossless beamsplitter has certain (complex-valued) probability amplitudes for sending an incoming photon into one of two possible directions. We use elementary laws of classical and

Hierarchy of quantum correlations using a linear beam splitter ...

In the present study, we investigate in detail the quantum correlations of continuous variable states through a beam splitter.

© 2026 DOLPHIN FIBER SYSTEMS - All rights reserved

the origin of anticorrelation for photon bunching on a beam splitter

Second-order anticorrelation in a beam splitter represents the origin of these phenomena and cannot be achieved classically.

The Quantum Regime Operation of Beam Splitters and Interference

The presence of quantum Rayleigh scattering, or spontaneous emission, inside a dielectric medium such as a beam splitter or an interferometric filter prevents a single photon from propagating

(PDF) Quantum physics and the beam splitter mystery

Non-polarizing beam-splitters (BSs) are the heart of most optical experiments and instruments (optical coherence tomography, holography, optical communication, quantum

Beamsplitters: A Guide for Designers | Optics

Alternately, other elements of the system can be designed to compensate for any aberrations introduced by the cube in a noncollimated beam. Cube beamsplitters consist of matched pairs of identical right

The origin of anticorrelation for photon bunching on a beam splitter

The Copenhagen interpretation has been long-lasted, whose core concepts are in the Heisenberg''s uncertainty principle and nonlocal correlation of EPR. The second-order anticorrelation

Hierarchy of quantum correlations using a linear beam splitter

The study of quantum correlations of continuous variable states with respect to a linear beam splitter in simple is interesting from perspective of basic physics and applications.

Hierarchy of quantum correlations using a linear beam splitter

Recently, a strict hierarchy relation of various kinds of quantum correlations in different passive and active quantum systems is shown, for instance, in a linear beam splitter , correlated ...

Quantum physics and the beam splitter mystery

ABSTRACT Optical lossless beam splitters are frequently encountered in fundamental physics experiments regarding the nature of light, including “which-way” determination of light particles, N.

Covering the Basics of Beamsplitters — Firebird Optics

© 2026 DOLPHIN FIBER SYSTEMS - All rights reserved

Polarizing Beamsplitter While standard non-polarizing beamsplitters divide light by wavelength, a polarizing beamsplitter will split the incident beam

How to Select a Beamsplitter

Power separating beamsplitters are used to split beams into two orthogonal paths, and can also combine portions of two different beams into one path to create a single, mixed beam. When a

Beam Splitter Input-Output Relations

Beam Splitter Input-Output Relations The beam splitter has played numerous roles in many aspects of optics. For example, in quantum information the beam splitter plays essential roles in teleportation,

Beam Splitter

The two beams of light return to the beam-splitter and are combined forming an image of the measured surface superimposed by an interference pattern on the image sensor array (camera). Usually a PSI

Why doesn''t a typical beam splitter cause a photon to decohere?

Experimentally, in a Mach-Zender interferometer we can fold light paths with a mirror while maintaining coherent interference, but passing either beam into the photocathode of a photodetector destroys

Related Video Reference

This video was associated with the source search result. Verify technical details against current product documentation and project requirements.

© 2026 DOLPHIN FIBER SYSTEMS - All rights reserved

Contact Us

Continue your research online:

Website: https://dolphinspreschool.co.za
Technical inquiry: https://dolphinspreschool.co.za/project-desk.html
African project coordination: Waterfall City, Midrand, Gauteng, South Africa

This document is for preliminary research. Verify specifications against current standards, product documentation and project requirements.

© 2026 DOLPHIN FIBER SYSTEMS - All rights reserved